arXiv:1304.2964v1 [gr-qc] 10 Apr 2013

The electric and magnetic spike solutions

Ernesto Nungesser [*][1,3] and Woei Chet Lim [†][2,3]

[1]Department of Mathematics, Royal Institute of Technology,
10044 Stockholm, Sweden
[2]Department of Mathematics, University of Waikato,
Private Bag 3105, Hamilton, New Zealand
[3]Max-Planck-Institute for Gravitational Physics,
Am Mühlenberg 1, 14476 Golm, Germany

April 11, 2013

Abstract

The aim of this paper is to use the existing relation between polarized Electrogowdy spacetimes and Gowdy spacetimes to find explicit solutions for electromagnetic spikes by a procedure which has been developed by one of the authors for gravitational spikes. We present new inhomogeneous solutions which we call the electric and the magnetic spike solutions.

1 Introduction

According to Belinskii, Khalatnikov and Lifshitz (BKL), a generic spacelike singularity is characterized by asymptotic locality. Asymptotically towards the initial singularity each spatial point evolves independently from its neighbors in an oscillatory manner that is represented by a sequence of Bianchi type I and II vacuum models. This picture has been recently refined. The starting point was when transformations found in [20] were used in [11] to find an explicit spike solution starting with a Kasner solution. This has then been developed in [11], [12] with the result that it seems that apart from local BKL behavior there also exist formation of spatial structures at and in the neighborhood of certain spatial surfaces that break asymptotic locality. Moreover the complete description of a generic spacelike singularity involves spike oscillations [5] which are described by sequences of Bianchi type I and certain inhomogeneous vacuum models. In [3] it has been shown that spikes occurring in a class of G_2 models lead to inhomogeneities that leave imprints on matter in the form of matter perturbations.

Here we present new solutions for the case of polarized Gowdy-symmetric spacetimes with an electromagnetic field. For this case the fundamental questions are on a firm ground. In [18] it was shown that the results of Ringström

---

[*] enun@kth.se
[†] wclim@waikato.ac.nz

concerning strong cosmic censorship for T^3-Gowdy spacetimes [22] could be translated to the case in consideration here since there exists a transformation of variables such that the system of partial differential equations is the same although the geometric interpretation is different. The aim of this paper is to use the existing relation between polarized/diagonal electromagnetic Gowdy spacetimes and Gowdy spacetimes to find the spike solutions for the electromagnetic case. The sign conventions of [19] are used. In particular, we use metric signature $-+++$ and geometrized units. We use the sign convention $\epsilon^{0123} > 0$ for the Levi-Civita tensor, but because we will switch to using a time variable that increases towards the past, the component ϵ^{0123} with respect to that coordinate system will be negative.

2 Basic equations and metric

We see that in order to compare the basic metric for polarized Gowdy spacetimes the relation between the coordinates (1) of [11] (with $Q = 0$) and (2) of [18] is obtained via $t = e^{-\tau}$ and $(\theta, x, y) = (x, y, z)$. The direction towards the initial singularity in this paper will thus corresponds to $\tau \to \infty$. In [18] a change of variables was performed in order to obtain the same system of PDE's mentioned. The change of variables is (9)–(10) of [18]:

$$P = 2\bar{P} - \tau \quad \lambda = 4\bar{\lambda} + 4\bar{P} - \tau, \quad \chi = \bar{Q}, \tag{1}$$

where $\bar{P}$, $\bar{\lambda}$ and $\bar{Q}$ refer to the vacuum solution. Our metric is:

$$ds^2 = -\mathrm{e}^{(\lambda-3\tau)/2} d\tau^2 + \mathrm{e}^{(\lambda+\tau)/2} dx^2 + \mathrm{e}^{P-\tau} dy^2 + \mathrm{e}^{-P-\tau} dz^2. \tag{2}$$

Note also that now we have electromagnetic radiation coming from a vector potential with only one nonzero component, namely $A_3 = \chi(x, \tau)$. The Einstein-Maxwell equations with P and χ are given by (19)–(24) of [17]:

$$\begin{aligned}
&P_{\tau\tau} - \mathrm{e}^{-2\tau} P_{xx} = 2(\chi_\tau^2 \mathrm{e}^{P+\tau} - \chi_x^2 \mathrm{e}^{P-\tau}) \\
&\chi_{\tau\tau} - \mathrm{e}^{-2\tau} \chi_{xx} = \mathrm{e}^{-2\tau} P_x \chi_x - (P_\tau + 1)\chi_\tau \\
&\lambda_\tau = -P_\tau^2 - \mathrm{e}^{-2\tau} P_x^2 - 4(\chi_\tau^2 \mathrm{e}^{P+\tau} + \chi_x^2 e^{P-\tau}) \\
&\lambda_x = -2P_\tau P_x - 8\mathrm{e}^{P+\tau} \chi_\tau \chi_x.
\end{aligned}$$

We will use the non-vanishing β-normalized variables [25].

$$\beta = \frac{1}{2} \mathrm{e}^{-\frac{\lambda - 3\tau}{4}}. \tag{3}$$

We take this opportunity to clarify the sign confusion in Eq. (9) of [11]. β and other kinematic variables are defined with respect to the future-pointing congruence. Therefore a positive β describes expansion towards the future (and contraction towards the past). We also take this opportunity to correct the error in the β expression in Eq. (2) of [12].

These β-normalized variables refer to the decompositions of orthonormal frame components of the 3-by-3 $\Sigma_{\alpha\beta}$ and $N_{\alpha\beta}$ matrices which in our case are

$$\Sigma_{\alpha\beta} = \begin{pmatrix} -2\Sigma_+ & 0 & 0 \\ 0 & \Sigma_+ + \sqrt{3}\Sigma_- & 0 \\ 0 & 0 & \Sigma_+ - \sqrt{3}\Sigma_- \end{pmatrix} \tag{4}$$

$$N_{\alpha\beta} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \sqrt{3}N_\times \\ 0 & \sqrt{3}N_\times & 0 \end{pmatrix}. \tag{5}$$

The β-normalized variables $\mathbf{Y} = (\Sigma_+, \Sigma_-, N_\times, \mathcal{E}, \mathcal{B})$, where $\mathcal{E}$ and $\mathcal{B}$ are β-normalized E^3 and B^2 orthonormal frame components of the electric and magnetic fields, are related to the partial derivatives of P, λ and χ as follows.

$$\mathbf{Y} = (\frac{1}{2} + \frac{1}{6}\lambda_\tau, -\frac{P_\tau}{\sqrt{3}}, -\frac{\mathrm{e}^{-\tau}P_x}{\sqrt{3}}, 2\chi_\tau e^{\frac{1}{2}(P+\tau)}, -2\chi_x e^{\frac{1}{2}(P-\tau)})$$

The evolution and constraint equations for the β-normalized variables are:

$$\begin{aligned}
\partial_t \Sigma_- &= \mathrm{e}^{-\tau}\partial_x(N_\times) + \frac{1}{2\sqrt{3}}(\mathcal{B}^2 - \mathcal{E}^2)] \\
\partial_t N_\times &= \mathrm{e}^{-\tau}\partial_x \Sigma_- - N_\times \\
\partial_t \mathcal{E} &= -\mathrm{e}^{-\tau}\partial_x \mathcal{B} + \frac{1}{2}\sqrt{3}N_\times \mathcal{B} + \frac{1}{2}(\sqrt{3}\Sigma_- - 1)\mathcal{E} \\
\partial_t \mathcal{B} &= -\mathrm{e}^{-\tau}\partial_x \mathcal{E} - \frac{1}{2}\sqrt{3}N_\times \mathcal{E} - \frac{1}{2}(\sqrt{3}\Sigma_- + 1)\mathcal{B} \\
\Sigma_+ &= \frac{1}{2}[1 - \Sigma_-^2 - N_\times^2 - \frac{1}{3}(\mathcal{E}^2 + \mathcal{B}^2)].
\end{aligned}$$

2.1 Hubble normalization

To visualize the dynamics we will use Hubble-normalized variables [26, 24], denoted with a superscript H. For the spatially homogeneous background dynamics, it is best to use the Hubble-normalized variables which are related to the β-normalized ones via

$$(\Sigma_+, \Sigma_-, N_\times, \mathcal{E}, \mathcal{B})^H = \frac{1}{1 - \Sigma_+}(\Sigma_+, \Sigma_-, N_\times, \mathcal{E}, \mathcal{B}), \tag{6}$$

and satisfy

$$\Sigma_+^{H2} + \Sigma_-^{H2} + N_\times^{H2} + \frac{1}{3}(\mathcal{E}^{H2} + \mathcal{B}^{H2}) = 1. \tag{7}$$

The Hubble-normalized energy density Ω of the electromagnetic field is given by

$$\Omega = \frac{1}{3}(\mathcal{E}^{H2} + \mathcal{B}^{H2}), \tag{8}$$

while the Hubble-normalized spatial curvature Ω_k is given by

$$\Omega_k = N_\times^{H2}. \tag{9}$$

2.2 Key transformations

The solution-generating transformations in polarized electromagnetic Gowdy spacetimes are

$$\mathrm{e}^{-\hat{P}/2} = \frac{\mathrm{e}^{-P/2}}{\chi^2 + \mathrm{e}^{-(P+\tau)}}, \quad \hat{\chi} = -\frac{\chi}{\chi^2 + \mathrm{e}^{-(P+\tau)}}, \tag{10}$$

and

$$\hat{P} = -P, \quad \hat{\chi}_\tau = -\mathrm{e}^{P-\tau}\chi_x, \quad \hat{\chi}_x = -\mathrm{e}^{P+\tau}\chi_\tau. \tag{11}$$

These transformations correspond to transformations (6) and (7) in [11], but curiously the role of solution-generating transformation has switched. Transformation (6) in [11] is merely a frame rotation, but here transformation (10) is a solution-generating transformation. On the other hand, transformation (7) in [11] is a solution-generating transformation, but here transformation (11) is merely a 90-degree duality rotation for the electromagnetic field (see e.g. [14] about duality rotation) followed by a switch of the coordinates y and z. Transformation (11), like transformation (7) in [11], is also very simple when expressed in β-normalized variables:

$$(\hat{\Sigma}_-, \hat{N}_\times, \hat{\mathcal{E}}, \hat{\mathcal{B}}) = (-\Sigma_-, -N_\times, \mathcal{B}, \mathcal{E}). \tag{12}$$

3 Explicit solutions

We now apply the vacuum-to-electromagnetic transformation (1) to the explicit solutions in Section 4 of [11].

3.1 Homogeneous solutions

3.1.1 Reparameterized Kasner solution

Applying the transformation (1) to the Kasner seed solution (17) of [11] yields

$$P = v\tau + 2P_0, \quad \chi = \chi_0, \quad \lambda = -v^2\tau + 4(\lambda_0 + P_0), \tag{13}$$

where $v = 2w-1$ and P_0, χ_0 and λ_0 are arbitrary constants. The result is trivial – this solution is just a re-parametrization of the Kasner solution. Nevertheless, we will use this parametrization of the Kasner solution in figure 3 later.

The β-normalized variables $\mathbf{Y_K}$ have the same form as (18) of [11]:

$$\mathbf{Y_K} = (\frac{1}{2} - \frac{1}{6}v^2, -\frac{v}{\sqrt{3}}, 0, 0, 0). \tag{14}$$

3.1.2 Electric Rosen solution

Applying (1) to the rotated Kasner solution (22) of [11] with the simplifying choice $P_0 + \ln\chi_0 = 0$ yields

$$\begin{aligned}
P &= -\tau - 2\ln\operatorname{sech} w\tau + 2\ln 2\chi_0 \\
\chi &= -\frac{1}{2\chi_0}(1 + \tanh w\tau) \\
\lambda &= -(4w^2+1)\tau - 4\ln\operatorname{sech} w\tau + 4(\lambda_0 + \ln 2\chi_0).
\end{aligned}$$

In terms of the β-normalized variables,

$$\mathbf{Y_{ER}} = (\frac{1}{3} + \frac{2}{3}w(\tanh w\tau - w), \frac{1}{\sqrt{3}}(1 - 2w\tanh w\tau), 0, -2w\,\mathrm{sech}\, w\tau, 0). \quad (15)$$

This describes an electric Bianchi I model. Up to the arbitrary constant and choosing $\chi_0 = 1$, $\xi = e^{-\tau}$ we have exactly the same χ as described in [2] with a minus sign where now $w = \frac{\alpha_0}{2}$ and $\gamma_0 = \lambda_0$. This solution was first found by Gerald Rosen [23] and we will refer to it as the electric Rosen solution with subscript ER.

3.1.3 Magnetic Rosen solution

Applying (1) to the Taub solution (24)–(26) of [11] yields

$$\begin{aligned} P &= \tau + 2\ln\mathrm{sech}\, w\tau - 2\ln 2\chi_0 \\ \chi &= 2\chi_0 wx + \chi_1 \\ \lambda &= -(4w^2+1)\tau - 4\ln\mathrm{sech}\, w\tau + 4(\lambda_1 - \ln 2\chi_0). \end{aligned}$$

We assume now that $\chi_1 = 0$ for simplicity. In terms of the β-normalized variables,

$$\mathbf{Y_{MR}} = (\frac{1}{3} + \frac{2}{3}w(\tanh w\tau - w), \frac{1}{\sqrt{3}}(2w\tanh w\tau - 1), 0, 0, -2w\,\mathrm{sech}\, w\tau). \quad (16)$$

We will refer to it as the magnetic Rosen solution and use the subscript MR. It is sometimes called the pure magnetic or magnetovacuum solution since later it has been generalized to include, e.g. a perfect fluid. It is not surprising that this solution appears here instead of the Bianchi II vacuum solution since there is a natural correspondence between heteroclinic chains consisting of Bianchi type II solutions in the vacuum case and heteroclinic chains in the case with a magnetic field which include orbits corresponding to both solutions of the vacuum Einstein equations of Bianchi type II and solutions of the Einstein-Maxwell equations of Bianchi type I. For recent work on oscillatory singularities in Bianchi models with magnetic fields we refer to [10].

3.2 Inhomogeneous solutions

3.2.1 Electric spike solution

Applying (1) to the rotated Taub solution (28)–(30) of [11] [1] and defining $f = xwe^{\tau}\,\mathrm{sech}\, w\tau$, $s = \frac{2f}{f^2+1}$, $c = \frac{f^2-1}{f^2+1}$ (note that $c^2 + s^2 = 1$), we obtain a new, inhomogeneous solution:

$$P = -3\tau - 2\ln\mathrm{sech}\, w\tau + 2\ln(f^2+1) + 2\ln 2\chi_0 \quad (17)$$

$$\chi = -\frac{1}{2\chi_0}\frac{f^2}{(f^2+1)xw} \quad (18)$$

$$\lambda = -(4w^2+9)\tau - 12\ln\mathrm{sech}\, w\tau + 4\ln(f^2+1) + 4(\lambda_1 + \ln 2\chi_0). \quad (19)$$

---

[1] As pointed out on page 14 of [12], the third minus sign in equation (28) of [11] should be a plus sign, and the factor 4 in equation (34) of [11] should not be there.

We will refer to it as the electric spike solution, because the spike occurs at $x = 0$, where the electric field is zero. Its β-normalized variables are given by

$$\mathbf{Y}_{\mathbf{ES}} = [-\frac{1}{3}(1+2w^2-2c+\sqrt{3}(c-2)\bar{\Sigma}_-), \frac{1}{\sqrt{3}}+c\bar{\Sigma}_-, s\frac{\mathcal{B}_{\mathrm{MR}}}{\sqrt{3}}, s\sqrt{3}\bar{\Sigma}_-, c\mathcal{B}_{\mathrm{MR}}] \quad (20)$$

where we have denoted $\bar{\Sigma}_- = (\Sigma_-)_{\mathrm{MR}} - \frac{1}{\sqrt{3}}$, and the subscripts MR and ES refer to the magnetic Rosen and the electric spike solution.

3.2.2 Magnetic spike solution

Applying (1) to the spike solution (33)–(35) of [11] yields

$$P = 3\tau + 2\ln\operatorname{sech} w\tau - 2\ln(f^2+1) - 2\ln 2\chi_0 \quad (21)$$

$$\chi = -\chi_0 w[\mathrm{e}^{-2\tau} + 2(w\tanh w\tau - 1)x^2] + \chi_2 \quad (22)$$

$$\lambda = -(4w^2+9)\tau - 12\ln\operatorname{sech} w\tau + 4\ln(f^2+1) + 4(\lambda_2 - \ln 2\chi_0). \quad (23)$$

We will refer to it as the magnetic spike solution and use the subscript MS . Its β-normalized variables are given by

$$\mathbf{Y}_{\mathbf{MS}} = [-\frac{1}{3}(1+2w^2-2c+\sqrt{3}(c-2)\bar{\Sigma}_-), -\frac{1}{\sqrt{3}}-c\bar{\Sigma}_-, -s\frac{\mathcal{B}_{\mathrm{MR}}}{\sqrt{3}}, c\mathcal{B}_{\mathrm{MR}}, s\sqrt{3}\bar{\Sigma}_-]. \quad (24)$$

Both the electric and magnetic spike solutions have nontrivial electric and magnetic fields, and the names ‘electric’ and ‘magnetic’ merely serve to differentiate the two. These solutions are new.

This completes the transformation of the explicit solutions in Section 4 of [11]. The solutions above can also be generated by starting with the Kasner solution (13) above and applying the transformations (10) and (11) successively.

4 Discussion



Figure 1. The electric and magnetic Rosen orbits projected on the $(\Sigma_+, \Sigma_-)^H$ plane.

We have already seen that the vacuum transformations have a different interpretation in the electromagnetic case. Now we proceed to visualize the dynamics of the different solutions and we compare the dynamics of the solutions found here corresponding to polarized Gowdy spacetimes with a electromagnetic field with the ones found in [11] corresponding to Gowdy spacetimes.

The dynamics of the electric and magnetic Rosen solutions can be described by their orbits in the state space. When projected on the $(\Sigma_+, \Sigma_-)^H$ plane, these orbits form straight lines emanating from one corner of a triangle superscribing the Kasner circle (see figure 1). The two transition sets combine to describe the dynamics during an electromagnetic equivalence of a Kasner era, which consists of long Kasner epochs (described by the Kasner equilibrium points), punctuated by brief periods of transitions. Either the electric or the magnetic component becomes significant during these periods of transitions. Compare with Figure 5 of [11]. The coincidence of the projected Rosen orbits with the projected Taub orbits compels one to compare the Rosen solutions with the Taub solution. Consider the Taub solution (24)–(26) of [11] with a particular value for w (call it w_{T}) and the corresponding magnetic Rosen solution whose orbit starts and ends at the same Kasner points as this Taub orbit. Then one finds that the w-parameter for the magnetic Rosen solution takes the value $w_{\mathrm{MR}} = w_{\mathrm{T}}/2$. Consequently the rate of change for the magnetic Rosen transition from one Kasner point to the next is only one half of that for the Taub transition. The implication of this is that in more general models where



Figure 2. Orbits of the spike solution projected on the $(\Sigma_+, \Sigma_-)^H$ plane. Orbits are colored red along the spike worldline $x = 0$, blue along $x = 1000$, and magenta along small values of x. $w = 0.2$, 0.5, 1, 1.5, 2, 3 respectively.

both modes are present, gravitationally-driven Taub transitions would dominate electromagnetically-driven Rosen transitions towards the singularity, and that the Hubble-normalized energy density of the electromagnetic field would tend to zero towards the singularity, with the caveat that this occurs almost everywhere, except possibly at some 'spiky' worldlines where the Taub mode has a local zero. Along these spiky worldlines, whether (vacuum) spike transitions would dominate Rosen transitions is unknown. In figure 2 we visualise the orbits of the magnetic spike solution (21)–(23) along various worldlines $x = \text{const}$, projected on the $(\Sigma_+, \Sigma_-)^H$ plane. The projected orbits of the electric spike solution (17)–(19) differ only in the sign of Σ_-. Along worldlines far away from the spike worldline, the orbits approximate the electric and magnetic Rosen orbits. Along the spike worldline, the projected orbit is a straight line. For the case $|w| \geq 1$, all these orbits end at the same Kasner points. For the case $0 < |w| < 1$, the orbit along the spike worldline ends at a different Kasner point from all others. Compare with Figure 6 of [11]. In figure 3 we visualise two families of the solutions (one with $w = 0.75$ and the other with $w = 1.25$). Note that the Kasner solution plotted here uses the parametrization in (13). Compare with Figure 9 of [11].



Figure 3. Two families of orbits with $w = 0.75$ and 1.25. The Kasner seed is indicated by a black *, the electric Rosen orbit in light blue, the magnetic Rosen orbit in dark blue, orbits of the electric spike solution in magenta, and orbits of the magnetic spike solution in red.

We now compare the orbits of magnetic spike solution with the orbits of vacuum spike solution. Call the corresponding parameters w_{MS} and w_{S}. We set $w_{\text{S}} = 2w_{\text{MS}} + 1$ so that both solutions start at the same Kasner point. But the solutions will not end at the same Kasner points. See figure 4.

5 Outlook

With the transformation which connects vacuum Gowdy spacetimes with polarized Gowdy-symmetric spacetimes we have found new inhomogeneous solutions which we have called electric and magnetic spikes. The spatially homogeneous



Figure 4. Comparison of orbits for the magnetic spike solution with $w_{\mathrm{MS}} = 1.5$ and orbits for the vacuum spike solution with $w_{\mathrm{S}} = 4$. Both sets of orbits start at the same Kasner point but end at two different Kasner points.

electric and magnetic Rosen solutions which have appeared here are of Bianchi type I. The asymptotic states of Bianchi I with a magnetic field where studied in [7]. Magnetic fields are also compatible with Bianchi II, VI_0, VII_0 and III and these combinations have already been studied [8], [9], [27], [6], [4].

A natural generalization would be to treat the full Gowdy-symmetric spacetimes with an electromagnetic field, see [13] for an introduction. For this situation strong cosmic censorship has *not* been shown. However there exist some global existence results [15], [16] and a decay result towards the future of certain quantity [21]. This last result has been extended to (vacuum) Gowdy spacetimes in higher dimensions [1].

Acknowledgements

We thank Alan A. Coley, Alan D. Rendall and Claes Uggla for helpful comments and suggestions. This work was initiated when both authors were still at the Max-Planck-Institute for Gravitational Physics.

References

[1] A. Barbos. *Energy decay law in n-dimensional Gowdy spacetimes with torus topology.* PhD thesis. Freie Universität Berlin, 2010.

[2] M. Carmeli, C. Charach, and S. Malin. Survey of cosmological models with gravitational, scalar and electromagnetic waves. *Physics Reports*, 76, 2:79–156, 1981.

[3] A. A. Coley and W. C. Lim. Generating matter inhomogeneities in general relativity. *Phys. Rev. Lett.*, 108:191101, 2012.

[4] S. R. Crowe. *Dynamical Systems Analysis of Magnetic Bianchi III Cosmologies.* MMath Thesis, University of Waterloo, 1998.

[5] J. M. Heinzle, C. Uggla, and W. C. Lim. Spike oscillations. *Phys. Rev. D*, 86:104049, 2012.

[6] J. T. Horwood and J. Wainwright. Asymptotic Regimes of Magnetic Bianchi Cosmologies. *Gen. Rel. Grav.*, 36:799–823, 2004.

[7] V. G. LeBlanc. Asymptotic states of magnetic Bianchi I cosmologies. *Class. Quant. Grav.*, 14:2281–2301, 1997.

[8] V. G. LeBlanc. Bianchi II magnetic cosmologies. *Class. Quant. Grav.*, 15:1607–1626, 1998.

[9] V. G. LeBlanc, D. Kerr, and J. Wainwright. Asymptotic states of magnetic Bianchi VI_0 cosmologies. *Class. Quant. Grav.*, 12:513–541, 1995.

[10] S. Liebscher, A. D. Rendall, and S. B. Tchapnda. Oscillatory singularities in Bianchi models with magnetic fields. *arXiv:1207.2655 [gr-qc]*, 2012.

[11] W. C. Lim. New explicit spike solutions-non-local component of the generalized Mixmaster attractor. *Class. Quantum Grav.*, 25:045014, 2008.

[12] W. C. Lim, L. Andersson, D. Garfinkle, and F. Pretorius. Spikes in the Mixmaster regime of G_2 cosmologies. *Phys. Rev. D*, 79:123526, 2009.

[13] P. A. Mansfield. *Cosmological singularities in electrovacuum spacetimes with two-parameter spacelike isometry groups.* PhD thesis. Yale University, 1989.

[14] C. W. Misner, K. S. Thorne, and J. A. Wheeler. *Gravitation.* Freeman, 1973.

[15] M. Narita. On the existence of global solutions for T^3-Gowdy spacetimes with stringy matter. *Class. Quant. Grav.*, 19:6279–6288, 2002.

[16] M. Narita. Global existence problem in T^3-Gowdy symmetric IIB superstring cosmology. *Class. Quant. Grav.*, 20:4983–4994, 2003.

[17] E. Nungesser. *Strong cosmic censorship in polarized T^3-Gowdy symmetric spacetimes with a Maxwell field.* Diploma thesis. Free University, Berlin, 2008.

[18] E. Nungesser and A. D. Rendall. Strong cosmic censorship for solutions of the Einstein-Maxwell field equations with polarized Gowdy symmetry. *Class. Quant. Grav.*, 26:105019, 2009.

[19] A. D. Rendall. *Partial differential equations in general relativity.* Oxford University Press, Oxford, 2008.

[20] A. D. Rendall and M. Weaver. Manufacture of Gowdy spacetimes with spikes. *Class. Quant. Grav.*, 18:2959–2976, 2001.

[21] H. Ringström. On the T^3-Gowdy Symmetric Einstein-Maxwell Equations. *Annales Henri Poincare*, 7:1–20, 2006.

[22] H. Ringström. Strong cosmic censorship in T^3-Gowdy spacetimes. *Annals of Mathematics*, 170:1181–1240, 2009.

[23] G. Rosen. Symmetries of the Einstein-Maxwell Equations. *J. Math. Phys.*, 3:313–318, 1962.

[24] C. Uggla, H. van Elst, J. Wainwright, and G. F. R. Ellis. The past attractor in inhomogeneous cosmology. *Phys. Rev. D*, 68:103502, 2003.

[25] H. van Elst, C. Uggla, and J. Wainwright. Dynamical systems approach to G_2 cosmology. *Class. Quant. Grav.*, 19:51–82, 2002.

[26] J. Wainwright and L. Hsu. A dynamical systems approach to Bianchi cosmologies: orthogonal models of class A. *Class. Quant. Grav.*, 6:1409–1431, 1989.

[27] M. Weaver. Dynamics of magnetic Bianchi VI_0 cosmologies. *Class. Quant. Grav.*, 17:421–434, 2000.